

# Chapter 1. Nearshore Fish use of embayment, spit, and bluff shoreline habitats of the central and western Strait of Juan de Fuca

Anne Shaffer and Tyler Ritchie, Washington Department of Fish and Wildlife, 332 E. 5<sup>th</sup> Street  
Port Angeles, Washington 98362

## **Introduction**

This report provides a summary of beach seining results from March 2007-August 2008 for shorelines of embayment, spit, and bluff sites, collectively called 'shoreline sites', of nearshore central and western Strait of Juan de Fuca, and highlights of interannual long term monitoring results observed to date at two embayed shoreline sites. While this report provides a detailed discussion of results from this portion of the Strait nearshore, the reader is also referred to individual watershed chapters for Pysht, Twins, Salt Creek, as well as the Chinook genetic report, forage fish use, kelp snorkeling surveys, and nearshore water quality reports.

## **Methods and Materials.**

Seining was conducted using the large Puget Sound Ambient Monitoring Protocols (PSAMP) protocol nets (Figure 1) deployed by boat from shore. Embayment, spit, and bluff sites were intended for sampling from March 2007-March 2008. Sites were to be sampled at a minimum of weekly from March-September 2007, and then monthly from October 2007 to March 2008. Two embayment sites, Freshwater Bay and Crescent Bay, continue to be sampled for long term monitoring, and have been sampled monthly from March-September 2008. On average two seines were conducted at each site and sampling day. For each seine all fish were identified to the lowest taxa possible, counted, and 25 of each species were measured to nearest mm. When observed, five Chinook were sampled for genetic analysis (see genetic report for methods). We opportunistically collected up to 100 fish of each forage fish species for biomass evaluation. These fish were preserved in 70 percent ethyl alcohol, returned to the lab, weighed, measured, and assessed for sexual maturity.

Fish densities were calculated to fish per cubic meter x 1000. Species richness and diversity were calculated for each site in the study.

## Results

A total of 181 seines were sampled from all shoreline sites from March 2007-March 2008. Sixty three percent of these were conducted between March and September 2007. Chronically limited agency boat resources and challenging sampling conditions of swell and fog severely limited our ability to sample Dungeness Bluffs, Dungeness Spit, Ediz Hook, and Elwha Bluffs, which are accessed only by boat. These sites were therefore sampled only from May thru October 2007. Due to further constraints of run time only one seine was done at spit and bluff sites. Data are summarized in Appendix A and B.

The majority of seines were at embayed (131 seines) sites followed by bluff (29 seines) and spit (25 seines) sites Overall, shoreline sites had the highest percentage of total fish caught over the year sampling period. The one exception was Ediz Hook, which had the second lowest percent of fish collected of all sites sampled over the entire period (Table 1).

Forage fish, specifically smelt, anchovy, sand lance, and herring, were the most abundant fish collected along all the shoreline sites (Table 1), and comprised well over 40% of total fish collected during the year sampling period. Bluff and spits had the highest weekly averages of total smelt (the majority of which were surf smelt see appendix for individual species). Smelt did show strong seasonal variation in weekly densities for all life histories (adults, juveniles, and post larval stages Figure 3a-f).

Continued monitoring of Freshwater and Crescent bay indicates that there is strong interannual variation in smelt and lingcod density. Total smelt densities at each site decreased by over 80% in 2008 compared to 2007, and percent composition by life history shifted from adult smelt in 2007 to juvenile smelt in 2008. (Figure 3g-h; Table 2&3.) Length of mature adult smelt did not vary appreciably between site. Maturity and weight however did show site differences (Figure 3e). Adult herring were not as frequently encountered, and most adult herring were not sexually mature or gravid (Figure 3f). Only one site in the Dungeness drift cell was observed to have sexually mature herring throughout the 13 month study.

For salmonid use of the nearshore, salmon were present at all embayment, spit, and bluff sites. The Twins nearshore had the lowest density of salmon, with only two coho collected from the Twins nearshore in December 2007. The Elwha drift cell, including Freshwater Bay, Elwha Bluffs, and Ediz Hook, had the highest density of Chinook and coho salmon (Figure 4). Genetic analysis of juvenile Chinook salmon collected during this survey revealed that a significant proportion of juvenile Chinook were from ESA listed stocks of both Puget Sound and Columbia river (Shaffer et al in prep, See Chapter 6 of the nearshore assessment report; Figure 4.).

### **Discussion**

Comparing diversity of shoreline sites, embayments, spits, and bluffs, indicates both trends and differences in ecological function. All sites had significant seasonal differences in diversity. Within embayments, Crescent Beach, Twins, and Pysht all have higher diversity than Freshwater Bay, which is degraded due to sediment starvation from Elwha river dams. Comparing between geomorphic habitat types of shoreline, the embayed sites had consistently higher diversity than spits and bluffs. Finally, the Elwha drift cell embayed, spit, and bluff sites had consistently lower diversity than comparative shoreline sites within the comparative Dungeness drift cell (Figure 5).

It is also important to note that, while the Elwha drift cell appears to be functioning at a lower level ecologically it is supporting the highest density of salmon, and in particular, Chinook salmon. In the genetic portion of this study we documented the majority of Chinook found along the central and western Strait shoreline to be from a number of listed stocks, including from Elwha Dungeness and Columbia river stocks (see Chapter 6 this report). Mapping of forage fish spawn from the Elwha drift cell has revealed that only Freshwater Bay supports surf smelt spawning.

Eelgrass mapping of the Elwha and comparative nearshore indicates remnant eelgrass beds in the Ediz Hook area, and extremely high use of understory kelp beds by juvenile forage fish (Norris et al 2007). This juxtaposition of current ecological fitness, current fish density and presence of numerous stocks of ESA listed priority recovery salmon species and forage fish

spawning, leads us to conclude that the Elwha drift cell, including embayed, spit, and bluff sites, is a top priority for nearshore restoration.

Restoration of the Elwha drift cell will be complex and requires modeling analysis. The Elwha drift cell is defined as extending from the western edge of Freshwater Bay to the tip of Ediz Hook. Sediment processes are the dominant limiting factor for the Elwha nearshore, and have played a significant role in nearshore habitat form (Warrick et al 2008). Historically the feeder bluffs east of the river mouth provided over 70% of sediment to the Elwha littoral system (Randall et al 2004; Shaffer et al 2008). The Elwha river provided the rest. The feeder bluffs east of the river mouth of the Elwha have been bulkheaded for over 100 years. Feeder bluffs in the lower river have been actively feeding since the dams were installed (Draut et al 2008). We therefore hypothesize that nearshore sediment starvation, and concomitant functional impacts, increase in Elwha nearshore with distance from the river mouth. This is supported by the documentation of surf smelt spawning, which requires a fine sand gravel substrate, only along the Elwha drift cell in immediate proximity to the Elwha river mouth.

Upcoming dam removals, which will partially restore Elwha nearshore sediment delivery, will have two phases. The first is anticipated to deliver 10mcy of sediment to the nearshore within five years of dam removal (see Randle et al 2004; Stolnack and Naimann 2005; and Shaffer et al 2008). The second will be the post dam removal annual rate of sediment delivery, estimated to be approximately 200,00cy per year. Prior to shoreline armoring the sediment contribution from the feeder bluffs was estimated to be 70% of sediment to the Elwha drift cell. After shoreline armoring this contribution was estimated to have dropped to zero. Armoring will remain in place, so sediment contribution from feeder bluffs within the Elwha drift cell will continue to be disrupted. Every effort should therefore be made to optimize this high volume but short interval of upcoming sediment delivery to the Elwha bluffs and Ediz Hook.

Additional specific modeling to define sediment fate in the Elwha nearshore and anticipated habitat response, in particular for predicting the proportion and fate of grain size that will be suitable for forage fish spawning and eelgrass recolonization, will be needed if the restoration actions are to be the most effective and successful.

The high diversity and richness of the embayed shorelines, and Crescent Bay in particular, make it, Freshwater Bay, and the Twins shoreline top priorities for long term preservation. The majority of these properties are privately owned. Acquisition and conservation easements of Crescent Bay and Freshwater Bay are therefore the appropriate and top priority actions for shoreline restoration. Crescent Bay also supported the highest numbers of sand lance of any other nearshore areas within the study area, including Pysht, which is a documented sand lance spawning beach. These consistently high numbers of sand lance lead us to recommend Crescent Bay continue to be monitored for sand lance spawning.

The Twins shoreline is intriguing in that it has high diversity, but salmon use was not heavily documented in this study. This despite heavy salmon use of the watershed documented by Roni et al 2008. We theorize that salmon use of the Twins nearshore, both shoreline and lower river, is much different than other nearshore areas of the central and western Strait. Additional monitoring is recommended to understand salmonid use of the Twins nearshore. The high diversity and species richness of the area, and high smelt use for both spawning and migration, leads us to recommend the Twins as a priority for restoration and acquisition. The Twins is a mixture of private and public ownership. Acquisition/conservation easement of private properties is a priority.

The Pysht shoreline also exhibited high species diversity and richness, as well as salmon use, including both Puget Sound and Columbia river ESA listed stocks. This area is the site of a large conservation easement. Every effort should be made to keep the Pysht shoreline intact. All development of this shoreline should be avoided.

Finally, preliminary results from ongoing long term sampling of two of the embayed sites indicates a large seasonal and interannual variation in fish use of the nearshore central Strait, with large differences in both total numbers of fish, and fish distribution in the nearshore. For example, between 30-60% fewer smelt were observed in Crescent and Freshwater Bay shoreline sites in 2008 relative to 2007 sampling, and the life history composition of smelt changed dramatically with year. This strong interannual variation may reflect a cyclic nature for smelt use

of the Strait, and partially explain large interannual differences in spawn that is characteristic of Strait smelt spawning beaches. Sand lance also had strong interannual differences in density. For both years, Crescent Bay had far more sand lance than Freshwater Bay. Also, in Crescent Bay 2008 we observed a strong and sustained recruitment of greenlings, primarily lingcod. This was not observed in 2007, but noted by WDFW biologists that worked in this area in the early 1980's (Doty, WDFW, pers comm.). This strong interannual variation makes it necessary to continue long term sampling of the nearshore central Strait so that we can clearly define current fish use of the nearshore, as well as changes in fish use that may occur due to restoration actions.

## References

- Norris, J, I. Ward, A. Shaffer and C. Lear 2007. Eelgrass mapping of the Elwha Nearshore. in Proceedings, Puget Sound Georgia Basin Conference , Puget Sound Water Quality Authority, Olympia Washington
- Randle, T. J., J. Bountry, B. Jackson, and G. Smille. 2004. Elwha River restoration project draft sediment management and monitoring plan, based on recommendations of the Elwha River Physical Processes Monitoring Workshop, 13-17 August, 2001, Port Angeles, WA.
- U.S. Department of the Interior, Bureau of Reclamation, Denver, CO and National Park Service, Fort Collins, CO.
- Roni, P. R. Holland, T. Bennett, G. Pess and R. Moses. 2008. Straits intensively monitored watershed contract report: results of FY07 PIT tagging on east and west Twin rivers. Northwest Fisheries Science Center, NMFS, Seattle, Washington 98112.
- Shaffer J.A., P. Crain, and Todd Kassler in prep. Juvenile Chinook use of the nearshore central and western Strait of Juan de Fuca. Washington department of fish and wildlife, Habitat program, science division.

Shaffer, J.A, P. Crain, B. Winter, M. McHenry, C. Lear and T. Randle. 2008. Nearshore Restoration of the Elwha River Through Removal of the Elwha and Glines Canyon Dams: An Overview. Northwest Science. 82 (Special Issue).

Stolnack, S. and R. Naiman. 2005. Summary of Research and Education Activities in the Elwha River Watershed and Adjacent Coastal Zone. University of Washington, Seattle.

U. S. Army Corps of Engineers. 1971. Report on survey of Ediz Hook for beach erosion and related purposes. Part II. Seattle District Army Corps of Engineers, Seattle, WA.


Warrick, J. A., G. R. Cochrane, Y. Sagy, and G. Gelfenbaum. 2008. *Nearshore substrate and morphology offshore of the Elwha River*. Northwest Science 82 (Special Issue):153-163.

**Acknowledgements.** The nearshore assessment project is sponsored by the North Olympic Peninsula Lead Entity and funded by the Salmon Recovery Funding Board (SRFB/RCO). Cheryl Baumann is the NOPE coordinator, Tara Galuska is the RCE project manager. The Clallam Marine Resources Committee (MRC) provided important funding for college interns including Jesse Charles, Chris DeSisto, Bryan Hara, Erica Hirsh, John Kascirk, Mario Laungayan, Romy Laungayan, Ross McDorman, Tiffany Nabors, Sean Oden, Rebecca Paradis, Jacob Ray, Tyler Ritchie, Melanie Roed, Justin Rondeau, Trista Simmons, Ben Warren, Karen Wilkie, Eric Wood, and Steve Wiyall from WWU, REU, and Peninsula College supervisors were Mr. Jack Ganzhorn and Dr. Dwight Barry of Peninsula College. Capt. Winn Miller and Phil Henry provided boat support and good will. Jenna Schilke, formerly with WDFW, supervised much of the field work. Dr. Jon Warrick, USGS, provided methods and the camera for photo analysis. The Clallam MRC continues to support student internships financially. Cathy Lear, Clallam County, is senior staff to the MRC. Dan Penttila, WDFW, provided report review.

REU is supported by the National Science Foundation's Research Experience for Undergraduates program under Grant DBI-0452328.

Chuck and Neva Novak and Ben and Irene Palzer continue to provide landowner access to the private beach at Crescent Bay. Malcolm Dudley has provided invaluable private land owner support and access. Pam Lowry has provided access to forage fish spawn sampling

locations in Freshwater Bay. Pam Sangunetti and Kevin Ryan (USFWS) provided invaluable partnership and field support. Charlie and Kendra Parks, Port Angeles High School, and Andy Stevenson, provided field support and good will. Thank you all.




 Shaffer et al, 2007  
WOPW


Figure 1. Strait nearshore assessment seine design provided by Christensen nets.

Table 1. Results. Percent of total fish and dominant species at each site, march 2007-march 2008.

Site	Total seines	Total Fish	% of all fish	Chinook	Coho	Chum	Smelt (adult = >120)	Smelt (juv = 50-120)	Smelt (pl = <50)	Herring (juv = 50-120)	Northern Anchovy	Sand lance (juv = 50-120)	3- Spine stickle back	Shiner perch	Stag horn sculpin	Cottids	Cutthroat	Prickley Sculpin
Crescent Bay	41	11563	6	2	2	2	<b>42</b>	<b>14</b>	0	4	0	<b>21</b>	1	1	3	0	0	0
Dungeness Bluffs	16	24960	13	0	0	0	1	6	<b>90</b>	2	0	0	0	0	0	0	0	0
Dungeness Spit	12	5049	3	0	0	6	4	<b>54</b>	<b>14</b>	<b>11</b>	0	8	0	0	0	0	0	0
Ediz Hook	9	3101	2	6	0	3	5	<b>20</b>	1	<b>51</b>	0	<b>12</b>	0	0	0	0	0	0
Elwha Bluffs	13	5518	3	10	1	9	1	<b>43</b>	<b>15</b>	<b>20</b>	0	1	0	0	0	0	0	0
Elwha W Estuary	52	16258	9	<b>12</b>	2	1	0	0	0	0	0	0	<b>70</b>	0	4	5	0	1
Freshwater Bay	28	12259	6	4	0	1	<b>14</b>	6	9	<b>56</b>	1	7	0	1	0	0	0	0
Pysht Shoreline	30	31720	17	0	0	0	<b>11</b>	<b>79</b>	0	2	0	0	0	0	3	0	0	0
Pysht Main Channel	44	9553	5	0	1	1	0	7	0	0	0	0	1	0	7	<b>79</b>	0	0
Pysht Side Channels	36	15586	8	0	0	0	0	0	0	0	0	0	<b>49</b>	<b>26</b>	<b>19</b>	3	0	0
Salt Creek Main Channel	66	18731	10	0	2	3	0	0	0	0	0	0	5	<b>68</b>	<b>14</b>	6	0	0
Salt Creek Side Channel	52	13947	7	0	1	2	0	0	0	0	0	0	<b>12</b>	<b>54</b>	<b>22</b>	7	0	0
W.Twins River	26	1477	1	0	0	1	0	0	0	0	0	0	0	0	5	<b>48</b>	6	<b>33</b>
W. Twins Shoreline	32	19810	10	0	0	0	<b>10</b>	<b>17</b>	4	<b>19</b>	<b>30</b>	0	0	0	0	20	0	0



percent of catch by site, fall winter 2007/2008  
(relative CPUE=39%)



- Crescent
- Freshwater Bay
- Pysht Shoreline
- Twins Shoreline
- Elwha Bluffs
- Dungeness Bluffs
- Dungeness Spit
- Ediz Hook
- Salt Creek Side Channel
- Elwha Side Channel
- Pysht Side Channels
- Pysht Main Channel
- Salt Creek Main Channel
- Twins Main




Figure 2 Percent of total fish collected by site by site for 2007-2008




Figure 3a. Average weekly total smelt density by geomorphic habitat type (fish per cubic meter) x1000.




Figure 3b. Average weekly post larval smelt density by geomorphic habitat type (fish per cubic meter) x1000.




Figure 3c. Average weekly juvenile smelt density by geomorphic habitat type (fish per cubic meter) x1000.




Figure 3d. Average weekly adult smelt density by geomorphic habitat type (fish per cubic meter) x1000.




Figure 3e. Weight, length and maturity of weighed adult smelt collected from shoreline sites. Maturity 1=Immature, 2=Starting, 3=Developing, 4=Maturing, 5=Mature, 6=Ripe, 7=Spent, 8=Recovering, 0=Juvenile






Figure 3f. Weight, length and maturity of weighed adult herring collected from shoreline sites. Maturity 1=Immature, 2=Starting, 3=Developing, 4=Maturing, 5=Mature, 6=Ripe, 7=Spent, 8=Recovering, 0=Juvenile






Figure 3g. Sand lance densities along shoreline sites.




Figure 3g. Long term monitoring results for average smelt densities Freshwater and Crescent Bays.




Figure 3h. Long term monitoring result, ling cod densities, Crescent Bay.




Figure 4. Chinook stock composition from embayment sites west of the Elwha river (Shaffer et al in prep, see accompanying Chapter 6 for additional details).

Table 2. Percent of total smelt for each year by life history. Total smelt % change based on CPUE.

Site	2007			2008			Percent Change in total smelt
	Adult	Juvenile	Post larval	Adult	Juvenile	Post larval	
Crescent	80	19	0	8	85	7	-32
Freshwater Bay	54	12	33	0	98	2	-60





Figure 4. Salmon density by site and species, March-September 2007.



Shannon Weiner Diversity (H') for Shoreline Sites







Figure 5. Ecological indices of Elwha and comparative sites by geomorphic habitat type.